

MEETLEARN.COM

Advanced Level Further Mathematics
Paper Three

Order Your Books at

www.shop.meetlearn.com

and receive them
right at your doorstep

JUNE 2001

- a) Two particles A and B have position vectors $(2 \sin 2t + 2 \cos 2t) \mathbf{j}$ m and $(2t + t^2) \mathbf{j}$ m respectively at time t seconds. Find the velocity of A relative to B when $t = 0$,

the acceleration of each particle when $t = \pi/4$

the value of t for which the velocities of A and B have the same magnitude,

the value of t for which the acceleration of A and B are parallel and in the same sense.

A particle of mass 2 kg is acted on by forces F_1 , F_2 and F_3 , where $F_1 = (4\mathbf{i} + 6\mathbf{j})$ N, $F_2 = (6\mathbf{i} + 8\mathbf{k})$ N, $F_3 = (2\mathbf{i} + 4\mathbf{k})$ N. Initially the particle is at rest at the point with position vector $(2\mathbf{i} - 3\mathbf{j} + \mathbf{k})$ m. Find the position vector and the magnitude of the momentum of the particle after 4 seconds. Find also, the work done on the particle in this time.

3. A particle p is projected from a point O of a horizontal plane π with velocity u at an angle α to the plane. Taking horizontal and vertical axes Ox and Oy respectively in the plane of the particle's path, show that the Cartesian equation of the particle's path is

$$Y = x \tan \alpha - gx^2 \sec^2 \alpha / 2u^2$$

The particle passes through the points $A(180, 30)$ and $B(240, 20)$. Find

- b) the time flight of P
- c) the range of P on A
- d) the greatest height of P above A

Take g as 10ms^{-2}


4. Two elastic spheres of masses 2m kg and 7m kg collide directly. Their velocities before impact are 6ms^{-1} and 3ms^{-1} respectively in the same direction and after impact they are $5/2 \text{ms}^{-1}$ and $u \text{ms}^{-1}$ respectively also in the same direction. Find the value of u and the coefficient of restitution between the spheres. Find, also, the velocities of the spheres after impact if they collide directly when travelling in opposite directions, the velocity of the lighter sphere being 7ms^{-1} that of the heavier sphere 2ms^{-1} . In this case determine

- 4. the loss in kinetic energy due to the impact,
- 5. the magnitude of the impulse experienced by the lighter sphere.

- ii) A car moving along a straight level road accelerates uniformly from rest until it has travelled a distance X m. It then, maintains a constant speed for 50 s and travels a further X m during this time. Finally it decelerates uniformly and comes to rest after travelling a further $X/2$ m. Find the total time for the journey.

(a) A particle P moves in a straight line. The speed $v \text{ m s}^{-1}$ of the particle at time t second is given by $V=kt^2 - ct$. When $t = 0$, the particle is at the origin and when $t = 1$, the particle is again at the origin and travelling with an acceleration of 4 m s^{-2} . Find the values of the constants k and c . Hence find the distance covered during the first 4 seconds of moving, the acceleration of P during the third second of motion

7. Two particles of masses $3m$ and $5m$ are attached to the ends of a light inextensible string of length a which passes through a smooth fixed ring at O. The lighter particle describes a horizontal circle about the heavier particle, which remains at rest, as centre. Show that the heavier particle is at a depth $3a/8$ below O, and that the time for complete revolution is $\pi\sqrt{3a/2g}$



7.

The figure above shows a block P of a mass 40 kg on a smooth horizontal table. The block P is connected by two light inextensible strings passing over two smooth pulleys to blocks Q and R of masses 10kg and 70kg respectively. The blocks hang freely and the system is released from rest with both strings taut and the hanging part vertical. Find

- (a) the magnitude of the acceleration of the system,
- (b) the tension in each string.

The block R falls a distance of 2m and is brought to rest by the floor. Calculate the farther distance that Q covers before momentarily coming to rest, given that it does not reach the pulley between it and block P.

[Take g as 10ms^{-2}]

8. i) An urn contains 10 red balls and 8 yellow balls. Balls are to be drawn from the urn at random, one after the other without replacement. Calculate the probability that

(a) the first two balls drawn will both be red balls

(b) the third ball drawn will be the first red ball

(c) the first ball will be green and the second will be yellow

(d) of the first two balls drawn, one will be green and the other will be yellow

(e) the last ball drawn will be the red

(ii) The events A and B are independent. The probability of event A occurring is $\frac{1}{5}$ and tip probability of event B occurring is $\frac{2}{5}$. Find the Probability of

(f) either A or B occurring,

(g) neither A nor B occurring ,

(h) One and only of the two events occurring